## Math 135, Calculus 1, Fall 2020

10-07: The Derivative, Graphs, and First Rules

Recall the **derivative function** f'(x) of a function f(x) is defined to be

- the slope of the tangent line
- the instantaneous velocity f(x + h) - f(x)
- $\lim_{h \to 0} \frac{f(x+h) f(x)}{h}$

A. Graphs of f'(x)

**Exercise 1.** Consider the function  $f(x) = x^2 - x - 2$ . We will construct the graph of the derivative function one point at a time.

- (a) Go to the website http://www.shodor.org/interactivate/activities/Derivate/.
- (b) Enter the function y = f(x) above. Use the tool to cacluate the slope of the tangent line at each of the points x = -1, 0, 1, 2, and 2.5. Enter these values in the table below:



(c) Now plot these points and connect them smoothly to estiamte a graph of f'(x).



(d) What do you think the formula for this graph is?

**Exercise 2.** Use the limit definition of the derivative to compute f'(x) for  $f(x) = x^2 - x - 2$ . Do your results match your results from Exercise 1?

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} =$$

**Exercise 3.** Desmos offers an interactive applet that gives a good visual and tactile experience of poroducing the derivative function point by point. Follow the link below and follow the instructions, using the function  $f(x) = \sin x$ .

https://www.desmos.com/calculator/jlcpl1spy2

In the space below, sketch the graphs of f(x) and f'(x). Do you recognize the graph of f'(x)?

2

## B. Differentiability

If the limit defining f'(a) exists, we say f is **differentiable** at x = a.

**Theorem 1.** If f(x) is differentiable at x = a, then it is also continuous there. However, the converse is **not** true: a function may be continuous at a point, but not differentiable there (e.g. f(x) = |x| is continuous at x = 0, but is not differentiable there).

What might go wrong? Any of the things that make a limit not exist. In paticular:

- (a) the left-hand limit might not equal the right-hand limit.
- (b) the limit might be infinite.

**Exercise 4.** Sketch the graph of a continuous function that is differentiable everywhere **except**:

- at *x* = 1 because the limit DNE as in (a) above (the derivative from the left does not equal the derivative from the right)
- at x = 4 because the limit DNE as in (b) above (the derivative is infinite)

**Notation 2.** The derivative of a function f(x) has a second notation, namely

$$f'(x) = \frac{dy}{dx}$$
  $f'(a) = \frac{dy}{dx}\Big|_{x=a}$ 

This alterantive notation  $\frac{dy}{dx}$  is read as "the derivative of *y* with respect to *x*", and is called **Leibniz notation**. It reminds us that the derivative is the slope:

$$m \equiv \frac{\Delta y}{\Delta x}$$
 so  $f'(x) = \frac{dy}{dx}$ 

Leibniz notation also helps to represent taking the derivatives as an operation: the symbol  $\frac{d}{dx}$  means "take the derivative with respect to x".

**Exercise 5.** Use the limit definition of the derivative to prove that the derivative of a line is its slope:

$$\frac{d}{dx}\left(mx+b\right) =$$